




UNDERSTANDING QUÉBEC HYDROPOWER

AMONG THE LOWEST GREENHOUSE GAS EMISSIONS OF ALL ELECTRICITY GENERATION OPTIONS

All forms of electricity generation emit greenhouse gases (GHG) over the course of their lifespan (construction, operation and decommissioning). For hydropower, GHG emissions are mainly carbon dioxide, and to a lesser extent, methane, resulting from decaying vegetation in flooded land. Based on a life cycle analysis, net GHG emissions from Québec hydropower are significantly lower than electricity generation from natural gas and coal, and on par with wind.¹

GHG EMISSIONS

Power generation options (g CO₂ eq.*/kWh)



METHANE IS NOT AN ISSUE IN QUÉBEC RESERVOIRS

Northern reservoirs emit little methane for two main reasons:

Location


Vegetation is very sparse in the **northern environments** where Hydro-Québec's reservoirs are located. Additionally, they are **far from agricultural or urban areas**, so run-off that reaches them is very low in organic matter and in nutrients. Less organic matter means that Québec reservoirs and lakes are less productive ecosystems than those in other regions.

Cold temperatures


Cold water contains **more dissolved oxygen** than warm water, leading to the formation of more carbon dioxide and less methane when organic matter decomposes. Some carbon in sediments decomposes to form methane due to low oxygen levels, but this will turn into carbon dioxide in the presence of oxygen in the water as it migrates to the surface. There is enough oxygen in one metre of the water column to oxidize the methane produced.

Temporary emissions

Hydro-Québec has been a pioneer in the study of greenhouse gas emissions from hydroelectric reservoirs. Our studies show that emissions peak immediately after reservoir creation, and **decline to natural lake levels within five to ten years.**²



¹ CIRAIG (2014). These results are similar to those published by the IPCC (2011); ² Tremblay et al. (2005)
* CO₂ = Carbon dioxide; CO₂ eq. = CO₂ equivalent



<1%

Methane's share of total emissions from Québec hydropower

Hydropower emissions are on a par with wind, **50 times lower** than natural gas, 70 times lower than coal, 5 times lower than solar

1993

Year Hydro-Québec's GHG research started

10

Years of study on the Eastmain reservoir in James Bay (before, during and after impoundment)

80

Number of experts involved in the Eastmain-1 GHG study

>120,000

Number of measurements taken

BIBLIOGRAPHY

Brothers, S., Y. T. Prairie & P. A. del Giorgio, 2012. Benthic and pelagic sources of carbon dioxide in boreal lakes and a young reservoir (Eastmain-1) in eastern Canada. *Global Biogeochem. Cycles* 26, doi:10.1029/2011GB004074.

Brothers, S., Y. T. Prairie, C. Teodoru & P. A. del Giorgio, 2011. Landscape heterogeneity influences CO₂ production in a young boreal reservoir. *Can. J. Fish. Aquat. Sci.* 69: 447-456.

CIRAIG (International Reference Centre for the Life Cycle of Products, Processes and Services). Comparing power generation options and electricity mixes. 2014. Various pagings 102 – [58] p.

Intergovernmental Panel on Climate Change (IPCC), 2011. Edenhofer, O., R. Pichs-Madruga, Y. Sokona, K. Seyboth, P. Matschoss, S. Kadner, T. Zwickel, P. Eickemeier, G. Hansen, S. Schloemer, C. von Stechow (Eds.) *Renewable Energy Sources and Climate Change Mitigation*, Cambridge University Press, 1075 pp. See Chapter 9, Figure 9.8.

Teodoru, C.R., J. Bastien, M.-C. Bonneville, P. del Giorgio, M. Demarty, M. Garneau, J.-F. Hélie, L. Pelletier, Y.T. Prairie, N. Roulet, I. Strachan & A. Tremblay, 2012. The Net Carbon Footprint of a Newly created Boreal Hydroelectric Reservoir. *Global Biogeochemical Cycles*, Vol 26, GB2016, DOI:10.1029/2011GB004187.

Teodoru, C.R., P. del Giorgio, Y. Prairie & M. Camire. 2009. pCO₂ dynamics in boreal streams of northern Quebec, Canada. *Global Biogeochemical Cycles* 23, GB2012, doi:10.1029/2008GB003404.

Teodoru, C.R., P. del Giorgio & Y.T. Prairie. 2010. Spatial heterogeneity of surface CO₂ fluxes in a newly created Eastmain-1 reservoir in northern Quebec, Canada. *Ecosystems*, 10.1007/s10021-010-9393-7.

Tranvik, L. J., J.A. Downing, J.B. Cotner, S. A. Loiselle, R. G. Striegl, T.J. Ballatore, P. Dillon, K. Finlay, K. Fortino, L.B. Knoll, P. L. Kortelainen, T. Kutser, S. Larsen, I. Laurion, D. M. Leech, S. L. McCallister, D. M. McKnight, J. M. Melack, E. Overholt, J. A. Porter, Y. Prairie, W.H. Renwick, F. Roland, B. S. Sherman, D. W. Schindler, S. Sobek, A. Tremblay, M. J. Vanni, E. von Wachenfeldt, E.D. Wachenfeldt & G. A. Weyhenmeyer. 2009. Lakes and Reservoirs as Regulators of Carbon Cycling and Climate. *Limnology and Oceanography*, 54 (6 part 2), 2298-2314.

Tremblay, A., J. Bastien, I. Strachan & M.-C. Bonneville, 2010. CO₂ and CH₄ fluxes at Eastmain-1 Reservoir (Quebec, Canada) using three different methods. *The International Journal on Hydropower & Dams*, Vol. 17, Issue 4, p. 78-83.

Tremblay, A., L. Varfalvy, C. Roehm & M. Garneau (Eds.). 2005. Greenhouse Gas Emissions: Fluxes and Processes, Hydroelectric Reservoirs and Natural Environments. *Environmental Science Series*, Springer, Berlin, Heidelberg, New York, 732 pages.

Youngil, K., N. T. Roulet, L. Changsheng, S. Frolking, I. B. Strachan, P. Changhui, C. R. Teodoru, Y. T. Prairie & A. Tremblay. 2016. Simulating carbon dioxide exchange in boreal ecosystems flooded by reservoirs. *Ecological Modelling*, 327:1-17.

For more information on hydropower generation and greenhouse gas emissions, please consult our Web site:

www.hydroquebec.com/sustainable-development/documentation-center/ghg-reservoir.html